

GRAVITATIONAL POTENTIAL ENERGY GROUP THEORETICALLY

Joachim Nzotungicimpaye*

Kigali Institute of Education, Department of Mathematics
P.O.Box 5039,Kigali-Rwanda
e-mail kimpaye @kie.ac.rw

February 7, 2008

Abstract

We show by symplectically realizing the one spatial Aristotle Lie group that the hamiltonian of the associated elementary system consist of a gravitational potential energy only.No kinetic term.

1 One spatial Aristotle Lie group A

We know that the Aristotle Lie group [1] $A(n)$, $1 \leq n \leq 3$ also called static group [2], is the space-time transformations group

$$x_0'^i = R_j^i x_0^j + x^i; t'_0 = t_0 + t \quad (1)$$

The multiplication law for this group is

$$(t, x^i, R_k^i)(t', x'^k, R_j^k) = (t + t', R_k^i x'^k + x^i, R_k^i R_j^k) \quad (2)$$

One sees that the Aristotle Lie group $A(n)$ is a direct product of the euclidean Lie group $E(n)$ and the time translation group. In this paper we interest us on the one spatial dimensional Lie group that we will simply denote by A . It is the one dimensional space-time translation Lie group. Its multiplication law is

$$(t, h)(t', h') = (t + t', h + h') \quad (3)$$

It is an abelian additive group . The central extension $\mathcal{G}$ of its Lie algebra is generated by P, E and M such that the non trivial Lie brackets are [3]

$$[P, E] = gM \quad (4)$$

g being a constant whose the physical interpretation will be given below. If the general element of the connected Lie group G generated by P, E and M is written as $\exp(\xi M)\exp(xP + tE)$, then the multiplication law for G is [3]

$$(\xi, t, h)(\xi', t', h') = (\xi + \xi' + ght', t + t', h + h') \quad (5)$$

*On absence from the University of Burundi

2 Coadjoint orbit of A

If we denote by $\mathcal{G}^*$ the dual of $\mathcal{G}$ and if the duality is defined by

$$\langle (m, e, p), (\delta\xi, \delta t, \delta x) \rangle = m\delta\xi + e\delta t + p\delta x \quad (6)$$

we find that the coadjoint action of the Aristotle Lie on $\mathcal{G}^*$ is

$$Ad_{(t,h)}^*(m, e, p) = (m, e - mgh, p + mgt) \quad (7)$$

The coadjoint orbit is then characterized by the invariant m . Let us denote it by $\mathcal{O}$. From (4) one verify that the symplectic form on $\mathcal{O}$ is

$$\sigma = dp \wedge dq \quad (8)$$

where $q = -\frac{e}{mg}$ and that the canonical action of the Aristotle Lie group is then

$$\phi_{(t,h)}(p, q) = (p + mgt, q + h) \quad (9)$$

The Aristotelian Lie algebra is then represented on $\mathcal{O}$ by the hamiltonian vector fields

$$\phi_*(P) = -\frac{\partial}{\partial q}; \phi_*(E) = -mg\frac{\partial}{\partial p} \quad (10)$$

One then can verify that the comomentum [4] $\lambda : \mathcal{A} \longrightarrow \mathcal{C}^\infty(\mathcal{O}, \mathfrak{K})$ is such that

$$\lambda(E) = -mgq; \lambda(P) = p \quad (11)$$

If we interpret g as the gravitational acceleration, then by (5) one can verify that the physical dimension for ξ is L^2T^{-1} where L and T are respectively the symbols for length and time. Moreover, if the physical dimension of (6) is that of an action (ML^2T^{-1}), then the physical dimensions for m, e and p are respectively mass, energy and linear momentum. From (9) one sees that the evolution law is given by

$$\phi_{(t,0)}(p(0), q(0)) = (p(0) + mgt, q(0)) \quad (12)$$

and that the generator of time translations is then represented by the dynamical vector field

$$X_H = \frac{\partial}{\partial t} - mg\frac{\partial}{\partial p} \quad (13)$$

from which one can verify that the hamiltonian is

$$H = mgq \quad (14)$$

We recognize in it the gravitational potential energy. From (13) one sees that the orbit is a static elementary particle of mass m , linear momentum p in the potential $V = mgq$ [5].

References

- [1] Souriau ,J.M., *Des particules aux ondes : quantification géométrique*, Symposium International Huygens principle,La Haye ,1990
- [2] Levy-Leblond, J.M.,*Journal of Mathematical Physics*, 9, 1605–, 1968
- [3] Hamermesh,M.,*Group theory and its applications to physical problems*, Addison Wesley, New York, 1962
- [4] Carinena ,J.F. ,*canonical group action*, ICTP report, IC/88/37
- [5] Any book on classical mechanics